

# Continuous Bose-Einstein condensation and superradiant clocks

**C. Beli<sup>1,2</sup>, S. Bennetts<sup>1,2</sup>, C.-C. Chen<sup>1,2,3</sup>, F. Famà<sup>1,2</sup>, R. González Escudero<sup>1,2,4</sup>,  
J. He<sup>1,2</sup>, B. Heizenreder<sup>1,2</sup>, G. A. Kazakov<sup>5</sup>, J. Minář<sup>6,2</sup>, B. Pasquiou<sup>1,2,7</sup>,  
S. Sarkar<sup>1,2</sup>, S. A. Schäffer<sup>1,2</sup>, F. Schreck<sup>\*1,2</sup>, M. Tang<sup>1,8</sup>, S. Zhou<sup>1,2</sup>**  
**and the iqClock consortium**

*1. Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, The Netherlands*

*2. QuSoft, Amsterdam, The Netherlands*

*3. NIST, Boulder, CO, USA*

*4. LaserLaB, Department of Physics and Astronomy, Vrije Universiteit, Amsterdam, The Netherlands*

*5. Atominstytut, The Vienna University of Technology, Austria*

*6. Institute for Theoretical Physics, Institute of Physics, University of Amsterdam, The Netherlands*

*7. Laboratoire de Physique des Lasers, CNRS, Université Sorbonne Paris Nord, Villetaneuse, France*

*8. Niels Bohr Institute, University of Copenhagen, Denmark*

Ultracold quantum gases are excellent platforms for quantum simulation and sensing. So far these gases have been produced using time-sequential cooling stages and after creation they unfortunately decay through unavoidable loss processes. This limits what can be done with them. For example it becomes impossible to extract a continuous-wave atom laser, which has promising applications for precision measurement through atom interferometry [1]. I will present how we achieved continuous Bose-Einstein condensation, creating BECs that persist in a steady-state for as long as we desire. Atom loss is compensated by feeding fresh atoms from a continuously replenished thermal source into the BEC by Bose-stimulated gain [2]. The only step missing to create the long-sought continuous-wave atom laser is the addition of a coherent atom outcoupling mechanism. In addition this BEC may give us access to interesting driven-dissipative quantum phenomena over unprecedented timescales. The techniques we developed to create the continuous source of thermal atoms are also nicely suited to tackle another challenge: the creation of a continuously operating superradiant clock [3]. These clocks promise to become more rugged and/or more short-term stable than traditional optical clocks, thereby opening new application areas. I will present how we are developing two types of superradiant clocks within the European Quantum Flagship consortium iqClock, the first operating on a kHz-wide transition of Sr [4] and the second on the mHz-narrow Sr clock transition.

## References

- [1] N. P. Robins, P. A. Altin, J. E. Debs, J. D. Close, *Phys. Rep.* **529**, 265 (2013)
- [2] C.-C. Chen, R. González Escudero, J. Minář, B. Pasquiou, S. Bennetts, F. Schreck, arXiv:2012.07605 (2020)
- [3] D. Meiser, J. Ye, D. R. Carlson, M. J. Holland, *Phys. Rev. Lett.* **102**, 163601 (2009)
- [4] H. Liu, S. B. Jäger, X. Yu, S. Touzard, A. Shankar, M. J. Holland, T. L. Nicholson, *Phys. Rev. Lett.* **125**, 253602 (2020)

---

\*Corresponding author: schreck@uva.nl